

Patient Education - Glycopyrrolate 0.8% Topical Cream

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 – Direct 385-900-1400, Toll Free 1-855-499-2556

Patient Education Insert

-Apply medication to affected area as prescribed

This medication was compounded by a pharmacy per prescriber request. For external use only. Wash hands before and after use unless the medication is for your hands. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain. Do not swallow and avoid contact with eyes. Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed. Doing any of these may increase the chance of undesirable side effects.

The “Do not use after” date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss an application, apply as soon as you remember. If it is close to the next application, skip the missed application. Do not double applications.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program

Website address: <https://www.fda.gov/Safety/MedWatch/default.htm> - Toll free number: 1-888-463-6332

Your medication contains the following active ingredients. Please review the following:

Glycopyrrolate - an anticholinergic agent. In this compounded preparation, the intended use is to decrease excessive sweating at the application sites as directed by your prescriber.

You should not use glycopyrrolate if you are allergic to it, or if you have:

a bladder obstruction or other urination problems a stomach or bowel obstruction (including paralytic ileus) severe constipation severe ulcerative colitis or toxic megacolon glaucoma myasthenia gravis or active bleeding with fast heartbeats, low blood pressure, shortness of breath, and cold hands or feet.

Tell your doctor if you have ever had:

an enlarged prostate liver or kidney disease heart problems or a heart rhythm disorder high blood pressure a stomach disorder such as ulcerative colitis, hiatal hernia, reflux disease, or slow digestion a colostomy or ileostomy a thyroid disorder or a nerve disorder.

It is not known whether glycopyrrolate will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

You should not breastfeed while using glycopyrrolate. Glycopyrrolate may slow breast milk production.

Do not give this medicine to a child without medical advice.

Get emergency medical help if you have signs of an allergic reaction: hives difficult breathing swelling of your face, lips, tongue, or throat.

Stop using glycopyrrolate and call your doctor at once if you have:

severe constipation, severe stomach pain and bloating diarrhea (especially if you have a colostomy or ileostomy) painful or difficult urination fast or pounding heartbeats, fluttering in your chest confusion, severe drowsiness eye pain, seeing halos around lights fever, shallow breathing, weak pulse, hot and red skin or

(in a child taking glycopyrrolate) dry diapers, fussiness, or excessive crying.

Common side effects may include:

constipation, nausea, vomiting, bloating drowsiness, dizziness, weakness, feeling nervous slow heartbeats sleep problems (insomnia) blurred vision, sensitivity to light dry mouth, decreased sense of taste decreased sweating, decreased urination impotence, sexual problems headache or rash.